

# ALLIGARE

## DICAMBA 4

### Specimen Label

Herbicide For Weed Control In Corn, Cotton, Sorghum, Soybean, Small Grains, Pasture, Hay, Rangeland, Farmstead (Non-Cropland), Fallow, Sugarcane, Asparagus, Turf And Grass

#### ACTIVE INGREDIENT:

Dimethylamine salt of dicamba (3,6-dichloro-0-anisic acid)\* .....50.2%

OTHER INGREDIENTS: .....49.8%

TOTAL: .....100.0%

\*This product contains 41.7% 3,6-dichloro-o-anisic acid (dicamba) or 4 pounds per gallon (480 g/L).

EPA Reg. No. 81927-55

## KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>IF IN EYES:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>Remove contact lenses, if present after the first 5 minutes, then continue rinsing eye.</li> <li>Call poison control center or doctor for treatment advice.</li> </ul>
<b>IF SWALLOWED:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF ON SKIN OR CLOTHING:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.	

#### Manufactured for:

Alligare, LLC • 13 N. 8<sup>th</sup> Street • Opelika, AL 36801

#### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING / AVISO

Causes substantial but temporary eye injury. Do not get in eyes, on skin, or on clothing. Harmful if swallowed. Avoid breathing spray mist. Harmful if absorbed through skin.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are made of any waterproof material.

All mixers, loaders, applicators and other handlers must wear:

- Long-sleeved shirt and long pants,
- Shoes plus socks,
- Goggles or faceshield, and
- Chemical-resistant gloves (e.g. barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, polyvinyl chloride (PVC) ≥14 mils, or Viton® ≥14 mils)

See engineering controls for additional requirements.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### ENGINEERING CONTROLS STATEMENTS

Pilots must use an enclosed cockpit in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)].

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Apply this product only as directed on label.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and green-houses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls worn over short-sleeve shirt and short pants
- Chemical-resistant footwear plus socks
- Chemical-resistant gloves (e.g. barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, natural rubber ≥14 mils, polyethylene, polyvinyl chloride (PVC) ≥14 mils, or Viton® ≥14 mils)
- Chemical-resistant headgear for overhead exposure
- Protective eyewear

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to the uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, in nurseries, in forests, or in greenhouses.

Do not enter or allow others to enter the treated areas until the spray has dried.

Before applying Alligare Dicamba 4, read all directions and precautions appearing on the container label and in this booklet. Failure to follow all directions and precautions may result in unsatisfactory weed control, crop injury, or illegal residues.

#### PRODUCT INFORMATION

The following directions apply to all uses of Alligare Dicamba 4. Additional precautions and restrictions will be found in each specific use section.

#### RESTRICTIONS:

Do not treat irrigation ditches or water used for crop irrigation or domestic uses.

Do not apply this product through any type of irrigation system.

Do not exceed the maximum single application rate of 2 pints (1.0 lb. a.i.) Alligare Dicamba 4 per application with no more than 2 applications per year.

#### RESISTANCE MANAGEMENT

Alligare Dicamba 4 is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to Alligare Dicamba 4 and other Group 4 herbicides. Weed species with acquired resistance to Group 4 may eventually dominate the weed population if Group 4 herbicides are used repeatedly in the same field or in successive years as the primary method of control for targeted species. This may result in partial or total loss of control of those species by Alligare Dicamba 4 or other Group 4 herbicides.

To delay herbicide resistance consider:

- Avoiding the consecutive use of Alligare Dicamba 4 or other target site of action Group 4 herbicides that have a similar target site of action, on the same weed species.
- Using tank-mixtures or premixes with herbicides from different target site of action Groups as long as the involved products are all registered for the same use, have different sites of action, and are both effective at the tank mix or prepack rate on the weed(s) of concern.
- Basing herbicide use on comprehensive IPM program.
- Monitoring treated weed populations for loss of field efficacy.
- Contacting your local extension specialist, certified crop advisors, and/or manufacturer for herbicide resistance management and/or integrated weed management recommendations for specific crops and resistant weed biotypes.

#### MIXING AND APPLICATION

UNLESS OTHERWISE SPECIFIED UNDER THE INDIVIDUAL USE HEADINGS OF THIS BOOKLET, THE FOLLOWING DIRECTIONS APPLY TO ALL CROP AND NON-CROP USES OF ALLIGARE DICAMBA 4. REFER TO INDIVIDUAL USE SECTIONS FOR ADDITIONAL PRECAUTIONS, RESTRICTIONS, APPLICATION RATES AND TIMINGS.

Alligare Dicamba 4 is a water-soluble formulation that can be applied using water or sprayable fluid fertilizer as the carrier. If a fluid fertilizer is to be used, a compatibility test (See COMPATIBILITY TEST) should be made prior to tank mixing.

Ground or aerial application equipment, which will give good spray coverage of weed foliage, should be used. HOWEVER, DO NOT USE AERIAL APPLICATION EQUIPMENT IF SPRAY PARTICLES CAN BE CARRIED BY WIND INTO AREAS WHERE SENSITIVE CROPS OR PLANTS ARE GROWING OR WHEN TEMPERATURE INVERSIONS EXIST.

Apply 3 to 50 gallons of diluted spray per treated acre when using ground application equipment or 1 to 10 gallons of diluted spray per treated acre (2 to 20 gallons of diluted spray per acre for preharvest uses) in a water-based carrier when using aerial application equipment. Use the higher level of the listed spray volumes when treating dense or tall vegetation. Use coarse sprays.

Select nozzles designed to produce minimal amounts of fine spray particles. Spray with nozzles as close to the weeds as is practical for good weed coverage.

To avoid uneven spray coverage, Alligare Dicamba 4 should not be applied during periods of gusty wind or when wind is in excess of 15 mph.

Avoid disturbing (e.g., cultivating or mowing) treated areas for at least 7 days following application.

#### BEST STEWARDSHIP PRACTICES

Alligare Dicamba 4 provides effective broadleaf weed and brush control when properly applied. Best stewardship practices in all mixing, loading, and application operations not only maximize weed control, but also protect ground and surface waters and minimize off-target movement.

This chemical is known to leach through the soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**GROUND AND SURFACE WATERS PROTECTION**

1) Point source contamination - To prevent point source contamination, do not mix, load this pesticide product within 50 feet of wells (including abandoned wells and drainage wells), sink holes, perennial or intermittent streams and rivers, and natural or impounded lakes and reservoirs. Do not apply pesticide product within 50 feet of wells. This setback does not apply to properly capped or plugged abandoned wells and does not apply to impervious pad or properly diked mixing/loading areas as described below.

Mixing, loading, rinsing, or washing operations performed within 50 feet of a well are allowed only when conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be on or move across the pad. The pad must be self-contained to prevent surface water flow over or from the pad. The pad capacity must be maintained at 110% that of the largest pesticide container or application equipment used on the pad and have sufficient capacity to contain all product spills, equipment or container leaks, equipment wash waters, and rainwater that may fall on the pad. The containment capacity does not apply to vehicles delivering pesticide shipments to the mixing/loading site. States may have in effect additional requirements regarding wellhead setbacks and operational containment.

Care must be taken when using this product to prevent: a) back siphoning into wells, b) spills or c) improper disposal of excess pesticide, spray mixtures or rinsates. Check valves or anti-siphoning devices must be used on all mixing equipment.

2) Movement by surface runoff or through soil - Do not apply under conditions which favor runoff. Do not apply to impervious substrates such as paved or highly compacted surfaces in areas with high potential for ground water contamination. Ground water contamination may occur in areas where soils are permeable or coarse and ground water is near the surface. Do not apply to soils classified as sand with less than 3% organic matter and where ground water depth is shallow (less than 8 feet in Arizona). To minimize the possibility of ground water contamination, carefully follow application rate recommendations as affected by soil type in the **Product Information** section of this label.

3) Movement by water erosion of treated soil - Do not apply or incorporate this product through any type of irrigation equipment nor by flood or furrow irrigation. Ensure treated areas have received at least one-half inch rainfall (or irrigation) before using tailwater for subsequent irrigation of other fields.

**SENSITIVE CROP PRECAUTIONS**

Alligare Dicamba 4 may cause injury to desirable trees and plants, particularly beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco, tomatoes, and other broadleaf plants when contacting their roots, stems or foliage. These plants are most sensitive to Alligare Dicamba 4 during their development or growing stage. FOLLOW THE PRECAUTIONS LISTED BELOW WHEN USING ALLIGARE DICAMBA 4.

- Do not treat areas where either possible downward movement into the soil or surface washing may cause contact of Alligare Dicamba 4 with the roots of desirable plants such as trees and shrubs.
- Avoid making applications when air currents may carry spray particles to areas where sensitive crops and plants are growing, or when temperature inversions exist. Do not spray near sensitive plants if wind is gusty or in excess of 5 mph and moving in the direction of adjacent sensitive crops. Leave an adequate buffer zone between area to be treated and sensitive plants. Coarse sprays are less likely to drift out of the target area than fine sprays.
- Use coarse sprays to avoid potential herbicide drift. Select nozzles, which are designed to produce minimal amounts of fine spray particles. Examples of nozzles designed to produce coarse sprays via ground application are Delavan Rain-drops, Spraying Systems XR flat fans, or large capacity flood nozzles such as D10, TK10, or greater capacity tips. Keep the spray pressure at or below 20 psi and the spray volume at or above 20 GPA, unless otherwise required by the manufacturer of drift-reducing nozzles. Consult your spray nozzle supplier concerning the choice of drift-reducing nozzles.
- Agriculturally approved drift-reducing additives may be used.
- Do not apply Alligare Dicamba 4 adjacent to sensitive crops when the temperature on the day of application is expected to exceed 85°F as drift is more likely to occur.
- To avoid injury to desirable plants, equipment used to apply Alligare Dicamba 4 should be thoroughly cleaned (See PROCEDURE FOR CLEANING SPRAY EQUIPMENT) before reusing to apply any other chemicals.

All crop uses of Alligare Dicamba 4 are intended for a normal growing interval between planting and harvest. No crop rotation restrictions exist if normal harvest of treated crop has occurred. If this interval is shortened, such as in cover crops that will be plowed under, do not follow up with the planting of a sensitive crop.

Crops growing under stress conditions such as drought, poor fertility, or foliar damage due to hail, wind or insects, can exhibit various injury symptoms that may be more pronounced if herbicides are applied.

Consult your local or state authorities for possible application restrictions and advice concerning these and other special local use situations. Tank mix recommendations are for use only in states where the tank mix product and application site are registered.

**BAND TREATMENTS**

Alligare Dicamba 4 may be applied as a band treatment. Use the formulas below to determine the appropriate rate and volume per treated acre.

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast RATE per treated acre} = \text{Band RATE per treated acre}$$

$$\frac{\text{Band width in inches}}{\text{Row width in inches}} \times \text{Broadcast VOLUME per treated acre} = \text{Band VOLUME per treated acre}$$

**COMPATIBILITY TEST**

Before mixing in the spray tank, it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (see following table).

**Amount of Herbicide to Add to One Pint of Spray Carrier (Assuming Volume is 25 Gallons per Acre)**

HERBICIDE FORMULATIONS	RATE PER ACRE	LEVEL TEASPOONS
Dry	1lb.	1 1/2
Liquid	1 pt.	1/2

If herbicide(s) do not ball-up or form flakes, sludge, gels, oily films or layers, or other precipitates, then the tested spray mix is compatible. Usually, incompatibility in any of the above-described forms will occur with 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent is recommended. Rerun the above COMPATIBILITY TEST with a suitable compatibility agent (1/4 teaspoon is equivalent to 2 pints per 100 gallons of fluid fertilizer).

**PROCEDURE FOR CLEANING SPRAY EQUIPMENT**

The steps listed below are suggested for thorough cleaning of spray equipment following applications of Alligare Dicamba 4 or tank mixes of Alligare Dicamba 4 or tank mixes of Alligare Dicamba 4 plus 2,4-D amine.

- Hose down thoroughly the inside as well as outside surfaces of equipment while filling the spray tank half full of water. Flush by operating sprayer until the system is purged of the rinse water.
- Fill tank with water while adding 1 quart of household ammonia for every 25 gallons of water. Operate the pump to circulate the ammonia solution through the sprayer system for 15 to 20 minutes and discharge a small amount of the ammonia solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
- Flush the solution out of the spray tank through the boom.
- Remove the nozzles and screens and flush the system with two full tanks of water.

The steps listed below are suggested for thorough cleaning of spray equipment used to apply Alligare Dicamba 4 as a tank mix with wettable powders (WP), emulsifiable concentrates (EC), or other types of water-dispersible formulations. Alligare Dicamba 4 tank mixes with water-dispersible formulations require the use of a water/detergent rinse.

- Complete step 1.
- Fill tank with water while adding 2 lbs. of detergent for every 40 gallons of water. Operate the pump to circulate the detergent solution through the sprayer system for 5 to 10 minutes and discharge a small amount of the solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
- Flush the detergent solution out of the spray tank through the boom.
- Repeat step 1, and follow with steps 2, 3 and 4.

**WEED LIST**

This is a general list of weeds which may be treated with Alligare Dicamba 4 in accordance with this label as specified under the rates and timing sections of the Individual Use headings. Proper usage of this product will give control or growth suppression of many ANNUAL, BIENNIAL, and PERENNIAL broadleaf weeds, and many WOODY brush and vine species including:

ANNUAL			
Amaranth, Spiny (Spiny Pigweed)	Evening Primrose, Cutleaf	Pennycress, Field (Fanweed, Frenchweed, Stinkweed)	Sicklepod
Aster, Slender	Fleabane, Annual	Pepperweed, Virginia (Peppergrass)	Sida, Prickly (Teaweed)
Bedstraw	Goosefoot, Nettleleaf	Pigweed, Prostrate	Smartweed, Green
Beggarweed, Florida	Henbit	Pigweed, Redroot (Carelessweed)	Smartweed, Pennsylvania
Broomweed, Common	Jimsonweed	Pigweed, Rough	Sneezeweeds, Bitter
Buckwheat, Wild	Knotweed	Pigweed, Smooth	Sowthistle, Annual
Buffalobur	Kochia	Pigweed, Triazine resistant (triazine resistant)	Sowthistle, Spiny
Burclover, California	Ladythumb	Purslane, Common	Spikeweed, Common
Burcucumber	Lambsquarters Common	Pursley, Florida	Spurge, Prostrate
Buttercup, Roughseed	Lambsquarters (triazine resistant)	Radish, Wild	Spurry, Corn
Carpeweed	Lettuce, Prickly	Ragweed, Common	Starbur, Bristly
Catchfly, Nightflowering	Mallow, Common	Ragweed, Giant (Buffaloweeds)	Sumpweed, Rough
Chamomile, Corn	Mallow, Venice	Ragweed, Lance-Leaf	Sunflower, Common (Wild)
Chickweed, Common	Mare's Tail (Horseweed)	Ragweed, bitter	Sunflower, Volunteer
Clovers (Annual)	Maryweed	Rubberweed, bitter	Thistle, Russian
Cockle, Corn	Morning-glory, Ivyleaf	Sesbania, Hemp	Thistle, Musk
Cockle, Cow	Morning-glory, Tall	Shepherdspurse	Thistle, Plumeless
Cocklebur, Common	Mustard, Tansy		
Croton, Tropic	Mustard, Wild		
Croton, Woolly	Mustard (Yellowtops)		
Daisy, English	Nightshade, Black		

BIENNIALS			
Burdock, Common	Geranium, Carolina	Plantain, Bracted	Thistle, Bull
Carrot, Wild (Queen Anne's Lace)	Gromwell	Ragwort, Tansy	Thistle, Milk
Cockle, White	Knapweed, Diffuse	Starthistle, Yellow	Thistle, Musk
Evening Primrose, Common	Knapweed, Spotted	Sweetclover	Thistle, Plumeless
	Mallow, Dwarf	Teasel	

PERENNIALS			
*Alfalfa	*Dock Broadleaf (Bitterdock)	Milkweed, Western	Sundrop, Halfshrub (Evening Primrose)
Artichoke, Jerusalem	*Dock, Curly	Milkweed, Whorled	Thistle, Canada
Aster, Spiny	Dogbane, Hemp	Nettle, Stinging	Toadflax, Dalmation
Bedstraw, Smooth	*Dogfennel (Cypressweed)	Nightshade, Silverleaf (White Hosenettle)	Tropical Soda Apple
Bindweed, Field	Fern, Bracken	Onion, Wild	Trumpet creeper (Buckvine)
Bindweed, Hedge	Garlic, Wild	*Plantain, Broadleaf	Vetch
Blueweed, Texas	Goldenrod, Canada	*Plantain, Buckhorn	Waterhemlock
*Bursage, (Bur Ragweed)	Goldenrod, Missouri	Pokeweed	Waterprimrose
Lakeweed, Povertyweed)	Goldenrod, Common	Ragweed, Western	Creeping
Buttercup, Tall	Hawkweed	Redvine	*Woodsorrel, Creeping
Campion, Bladder	Henbane, Black	Sericia Lespedeza	Common Yellow
Chickweed, Field	Horsenettle, Carolina	Smartweed, Swamp	Common Yellow
Chickweed (Mooseear, Canada)	Ironweed	Snakeweed, Broom	Common Yellow
Chicory	Knapweed, Black	*Sorrel, Red (Sheep Sorrel)	Common
*Clover, Hop	Knapweed, Russian	Sowthistle	Wormwood
*Dandelion, Common	Milkweed, Climbing	Sowthistle, Perennial	Louisiana
	Milkweed, Common	Spurge, Leafy	*Yankeeweed
	Milkweed, Honeyvine		Yarrow, Common

\*Noted perennials may be controlled using Alligare Dicamba 4 at rates lower than those recommended for other listed perennial weeds. (See application rates and timing sections in this label.)

WOODY			
Alder	*Dewberry	Locust, Black	Sagebrush, Fringed
Ash	*Dogwood	Maple	Sassafras
Aspen	Elm	Mesquite	Serviceberry
Basswood	Grape	Oak	Spicebush
Beech	*Hawthorn (Thornapple)	Oak, Poison	Spruce
Birch	Hemlock	Olive, Russian	Sumac
*Blackberry	Hickory	Persimmon, Eastern	*Sweetgum
*Blackgum	Honeylocust	Pine	Sycamore
*Cedar	Honeysuckle	*Plum, Sand (Wild Plum)	Tarbrush
Cherry	Hornbeam	Poplar	Willow
Chinquapin	Huckleberry	Rabbitbrush	Witchhazel
Cottonwood	Huisache	*Redcedar, Eastern	*Yaupon
*Creosotebush	Ivy, Poison	*Rose, McCartney	*Yucca
Cucumbertree	Kudzu	*Rose, Multiflora	

\*Growth suppression

#### FIELD, SEED\*, POPCORN\* AND SILAGE CORN

**IMPORTANT:** Observe all restrictions, precautions, mixing and application instructions.

#### RESTRICTIONS:

- Do not apply Alligare Dicamba 4 to seed corn or popcorn without first verifying with your local seed corn company (supplier) the Dicamba selectivity on your inbred line or variety of popcorn. This precaution will help avoid potential injury of sensitive varieties.
- Alligare Dicamba 4 is not registered for use on sweet corn.
- Direct contact of Alligare Dicamba 4 with corn seed must be avoided. If corn seeds are less than 1 1/2 inches below the surface, delay application until corn has emerged.
- Up to 2 applications of Alligare Dicamba 4 may be made during a growing season. Do not exceed a total of 1 1/2 pints of Alligare Dicamba 4 per treated acre per crop year. Allow two weeks or more between applications of Alligare Dicamba 4. See appropriate section for rate information. For combination options or sequential treatments, refer to appropriate section.
- Do not use adjuvants containing penetrants such as petroleum-based oils after crop emergence or crop injury may result.
- Corn may be harvested or grazed for feed once the crop has reached the ensilage (milk) stage or later in maturity.

#### PRECAUTIONS:

- Applications of Alligare Dicamba 4 to corn during periods of rapid growth may result in temporary leaning. Corn will usually become erect within 3 to 7 days. Cultivation should be delayed until after corn is growing normally to avoid breakage.
- Agriculturally approved surfactants or sprayable fertilizers (1/2 to 1 gallon per acre of 28%, 30% or 32% urea ammonium nitrate or 2.5 pounds per acre spray grade ammonium sulfate) may be added to the spray mixture to improve postemergence weed control, particularly in dry growing conditions.
- Several synthetic pyrethroid insecticides are labeled for tank mix applications of dicamba. Refer to their label for specific directions, restrictions and precautions.

#### WEEDS CONTROLLED

Alligare Dicamba 4 will control many ANNUAL broadleaf weeds or give growth suppression of many PERENNIAL broadleaf weeds commonly found in corn. (Refer to the WEED LIST).

For best performance, make application when weeds have emerged and are actively growing.

Preemergence control of cocklebur, velvetleaf, and jimsonweed may be reduced if conditions such as low temperature or lack of soil moisture cause delayed or deep germination of weeds.

#### PREPLANT/PREEMERGENCE IN NO-TILLAGE CORN

Applications of Alligare Dicamba 4 may be made before, during, or after planting to emerged and actively growing broadleaf weeds. Apply Alligare Dicamba 4 at 1 pint per treated acre on medium or fine textured soils containing 2% or greater organic matter. Use 1/2 pint per treated acre on coarse textured soils (sand, sandy loam, and loamy sand) or medium and fine textured soils with less than 2% organic matter.

When planting into a legume sod (e.g., alfalfa or clover), apply Alligare Dicamba 4 after 4 to 6 inches of regrowth has occurred.

#### TANK MIX TREATMENTS FOR CORN

Alligare Dicamba 4 may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

Alligare Dicamba 4 Plus	Preplant/ Preemergent (No Tillage Corn)	RATES AND TIMINGS			Additional Directions
		Pre-emergent (Conventional or Reduced Tillage Corn)	Early Post-Emergent (All Tillage Systems)	Late Post-Emergent (All Tillage Systems)	
Accent® (nicosulfuron)	-	-	1/2-1 oz a.i./A	1/2-1 oz a.i./A (To improve spray coverage of weeds and reduce risk of corn injury, use drop pipes to direct spray beneath corn leaves when corn is greater than 8 inches tall)	Application may be made to emerged weeds before corn is greater than 24 inches tall. Use non-ionic surfactant at 0.25% (v/v) with this tank mixture.
Atrazine	1 1/4-2 lbs a.i./A	1 1/4-2 lbs a.i./A	1 1/4-2 lbs a.i./A Crop oil concentrates may be used with this mixture if corn is 5 inches or less in height.	1 1/4-2 lbs a.i./A Do not apply if corn is greater than 12 inches tall.	Application may be made before grasses are 1 1/2" tall. Follow all state and Federal restrictions pertaining to atrazine applications.
Beacon® (primisulfuron)	-	-	0.31-0.62 oz a.i./A	0.31-0.62 oz a.i./A (To improve spray coverage of weeds and reduce risk of corn injury, use drop pipes to direct spray beneath corn leaves when corn is greater than 8 inches tall)	Application may be made to emerged weeds when corn is 4 to 24 inches tall. Use non-ionic surfactant at 0.25% (v/v) with this tank mixture.
Metolachlor	1 1/2-3 lbs a.i./A	1 1/2-3 lbs a.i./A (Use only on fine or medium textured soils with 2 1/2% or greater organic matter.)	1 1/2-3 lbs a.i./A	-	Application may be made before grasses reach the 2 leaf stage and before corn is greater than 3 inches tall.
Frontier® (dimethenamid)	13-25 fl oz/A	13-25 fl oz/A (Use only on fine or medium textured soils with 2.5% or greater organic matter.)	13-25 fl oz/A	-	Application may be made up to 8 inch tall corn. This treatment must be combined with a herbicide that provides post-emergence control of grass weeds if they are greater than 1 inch tall at the time of application.

#### PREEMERGENCE IN CONVENTIONAL OR REDUCED TILLAGE CORN

Alligare Dicamba 4 may be applied after planting and prior to corn emergence. Application at 1 pint per treated acre may be made to medium or fine textured soils, which contain 2% or greater organic matter. DO NOT apply to coarse textured soils (sand, sandy loam, and loamy sand) until after crop emergence (see Early Postemergence uses below).

Preemergence application of Alligare Dicamba 4 does not require mechanical incorporation to become active. A shallow mechanical incorporation is recommended if application is not followed by adequate rainfall or sprinkler irrigation. Avoid tillage equipment (e.g., drags, harrows) which concentrates treated soil over seed furrow.

#### EARLY POSTEMERGENCE (ALL TILLAGE SYSTEMS)

##### (Spike through 8-inch tall corn)

Alligare Dicamba 4 at 1 pint per treated acre may be applied during the period from corn emergence through the five leaf stage or 8 inches tall, whichever comes first. Reduce the rate to 1/2 pint per treated acre if corn is growing on coarse textured soils (sand, sandy loam, and loamy sand). See LATE POSTEMERGENCE APPLICATIONS given below if the 6th true leaf is emerging from whorl or corn is greater than 8 inches tall.

#### LATE POSTEMERGENCE (ALL TILLAGE SYSTEMS)

##### (8 to 36 inch tall corn)

Application of Alligare Dicamba 4 at 1/2 pint per treated acre may be made from 8 to 36 inch tall corn or 15 days before tassel emergence, whichever comes first. For best performance, make applications when weeds are less than 3 inches tall.

Make directed spray application when (1) corn leaves prevent proper spray coverage; (2) sensitive crops are growing nearby; (3) tank mixing with 2,4-D. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

#### RESTRICTIONS:

DO NOT apply Alligare Dicamba 4 when soybeans are growing nearby if any of these conditions exist:

- corn is more than 24 inches tall
- soybeans are more than 10 inches tall
- soybeans have begun to bloom

#### OVERLAY (SEQUENTIAL) TREATMENTS

Alligare Dicamba 4 may be applied to ground previously treated with one or more of the following herbicides registered for use in corn:

acetochlor	glyphosate
alachlor (Lasso®, Lasso MT®, Partner®)	halosulfuron (Battalion®, Permit®, Lariat®)
atrazine	metolachlor
Broadstrike®	paraquat
butylate (Sutan®)	pendimethalin
dimethenamid (Frontier®)	propachlor (Ramrod®)
EPTC	simazine

Apply Alligare Dicamba 4 at 1/2 pint per treated acre to ground previously treated with full rates of Clarity or Marksman herbicides. Allow at least 2 weeks between applications.

READ AND FOLLOW LABEL DIRECTIONS FOR EACH OF THE ABOVE PRODUCTS.

Alligare Dicamba 4 Plus	RATES AND TIMINGS				Additional Directions
	Preplant/ Preemergent (No Tillage Corn)	Pre-emergent (Conventional or Reduced Tillage Corn)	Early Post-Emergent (All Tillage Systems)	Late Post-Emergent (All Tillage Systems)	
Frontier® 6.0 (dimethenamid)	16-32 fl oz/A	16-32 fl oz/A (Use only on fine or medium textured soils with 2.5% or greater organic matter.)	-	-	Application may be made up to 8 inch tall corn. This treatment must be combined with a herbicide that provides post-emergence control of grass weeds if they are greater than 1 inch tall at the time of application.
Paraquat	1/4-1 lb a.i./A	1/4-1 lb a.i./A	-	-	Application may be made to emerged weeds but prior to corn emergence.
Acetochlor	1 1/2-3 lbs a.i./A	1 1/2-3 lbs a.i./A (Use only on fine textured soils with greater than 2.5% organic matter)	-	-	Application should be made prior to corn emergence.
Lasso® (alachlor)	1 1/2-4 lbs a.i./A	1 1/2-4 lbs a.i./A (Use only on fine textured soils with greater than 2.5% organic matter.)	1 1/2-4 lbs a.i./A	-	Application may be made before grasses reach the 2 leaf stage and before corn is greater than 3 inches tall. If microencapsulated forms of alachlor are used (Lasso MT Partner), applications must be made prior to grass emergence.
Simazine	2.0-3.0 lbs a.i./A	2.0-3.0 lbs a.i./A	-	-	Application may be made prior to corn or weed emergence.
Pendimethalin	-	3/4-1 1/2 lbs a.i./A (Use only on fine or medium textured soils with 2 1/2% or greater organic matter.)	3/4-1 1/2 lbs a.i./A	-	Application may be made immediately after planting but prior to weed emergence. Corn should not be beyond the 2 leaf stage of growth.
Glyphosate	1.0-3.0 lbs a.i./A	1.0-3.0 lbs a.i./A	-	-	Application may be made to emerged weeds but prior to corn emergence.
Clopyralid	-	-	0.035-0.07 lb a.i./A	0.035-0.07 lb a.i./A	Application may be made any time after corn emergence through 24 inch tall corn. Use drop nozzles to direct spray after corn exceeds the 8 inch stage. Apply when the majority of the thistle plants have emerged and are at least 4 inches in height, but before bud stage. Use higher rates listed for stand reduction or larger thistle plants or heavier infestations. Lower rates listed may provide seasonal thistle suppression only.
Tough® (pyridate)	-	-	0.47 lb a.i./A	0.47 lb a.i./A	Application may be made to emerged, actively growing weeds. Directed applications are recommended when corn is large enough to prevent proper spray coverage.
2,4-D	1/4-1/2 lb a.i./A	1/4-1/2 lb a.i./A	Not recommended	1/8 lb a.i./A	Drop pipes are to be used when corn height is 8 inches or greater. Keeping the spray off the corn leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of weed foliage.

### COTTON EXCEPT CALIFORNIA

**IMPORTANT:** Observe all restrictions, precautions, mixing, and application instructions.

**PREPLANT APPLICATION:** Apply up to 8 fluid ounces of Alligare Dicamba 4 per acre to control emerged broadleaf weeds prior to planting cotton in conventional or conservation tillage systems.

For best performance, apply Alligare Dicamba 4 when weeds are in the 2 - 4 leaf stage and rosettes are less than 2" across.

#### RESTRICTIONS:

- Do not apply preplant to cotton west of the Rockies.
- Do not make Alligare Dicamba 4 preplant applications to geographic areas with average annual rainfall less than 25".
- If applying a spring preplant treatment following application of a fall preplant (postharvest) treatment, then the combination of both treatments may not exceed 2 pounds acid equivalent per acre.
- Following application of Alligare Dicamba 4 and a minimum accumulation of 1" of rainfall or overhead irrigation, a waiting interval of 21 days is required per 8 fluid ounces per acre or less. These intervals must be observed prior to planting cotton.

#### COTTON TANK MIXES

For control of grasses or additional broadleaf weeds, Alligare Dicamba 4 may be tank mixed with prometryn, paraquat, and glyphosate herbicides. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

#### SORGHUM (MILO)

**IMPORTANT:** Observe all restrictions, precautions, mixing, and application instructions.

#### RESTRICTIONS:

- Pre-Harvest Interval (PHI):** Grain sorghum (PHI): 30 days; Fodder (PHI): 30 days; Forage (PHI): 20 days
- Do not graze or feed treated sorghum forage or silage prior to mature grain stage. If sorghum is grown for pasture or hay, refer to the pasture use section of this label.
- Do not apply Alligare Dicamba 4 to sorghum grown for seed production.
- Make no more than one application per growing season.

#### PRECAUTION:

- Applications of Alligare Dicamba 4 to sorghum during periods of rapid growth may result in temporary leaning of plants or rolling of leaves. These effects are usually outgrown within 10 to 14 days.

#### WEEDS CONTROLLED

Alligare Dicamba 4, when applied at the listed rate for sorghum, will control many actively growing ANNUAL broadleaf weeds and will reduce competition from established PERENNIAL broadleaf weeds as well as control their seedlings. (Refer to WEED LIST).

#### RATES AND TIMINGS

Alligare Dicamba 4 may be applied to emerged and actively growing weeds at least 15 days prior to planting. Postemergence application of Alligare Dicamba 4 must be made after sorghum is in the spike stage (all sorghum emerged) but before sorghum is 15 inches tall. For best performance, make applications when sorghum is in the 3 to 5 leaf stage and weeds are small (less than 3 inches tall). Use drop pipes (drop nozzles) if sorghum is taller than 8 inches. Keeping the spray off the sorghum leaves and out of the whorl will reduce the likelihood of crop injury and improve spray coverage of weed foliage.

**Broadcast rate per treated acre:** 1/2 pint (1/4 lb. a.i.)

#### TANK MIX TREATMENTS

Alligare Dicamba 4 may be tank mixed with the products listed below. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

#### Alligare Dicamba 4 plus Atrazine:

For improved control of emerged, actively growing broadleaf weeds including triazine resistant species and added suppression of perennial broadleaf weeds, tank mix 1/2 pint Alligare Dicamba 4 with 0.5 to 1.25 lbs. a.i. atrazine per treated acre. For control of grasses (less than 1.5 inches tall), tank mix 1/2 pint Alligare Dicamba 4 with 2 lbs. a.i. atrazine per treated acre. For best performance and minimal crop injury, make application when sorghum is 3-8 inches tall and when broadleaf weeds are small (less than 6 inches tall). Application of atrazine must be made before sorghum is beyond 12 inches tall. The atrazine rate will depend upon soil texture and length of residual weed control desired. Follow all state and Federal restrictions pertaining to atrazine applications.

#### Alligare Dicamba 4 plus bromoxynil:

For improved control of broadleaf weeds, tank mix 1/2 pint Alligare Dicamba 4 with 1 - 1 1/2 pints bromoxynil herbicide per treated acre. Make application at 4 leaf to 15-inch tall sorghum. Use drop nozzles to direct spray beneath sorghum leaves when sorghum is greater than 8 inches tall.

#### OVERLAY (SEQUENTIAL) TREATMENTS

Alligare Dicamba 4 may be applied to ground previously treated with one or more of the following herbicides:

Herbicide	Maximum Rate Per Treated Acre (lbs. a.i.)
alachlor (Lasso®) (Screen® - treated seed)	4
atrazine <sup>1</sup>	2.5
metolachlor	2.5
propachlor (Ramrod®)	5

<sup>1</sup> Maximum use rate for atrazine is determined by soil type, tillage practices used, surface residue, and state or local restrictions. Follow the more restrictive requirements when determining the maximum use rate for atrazine.

**PREHARVEST USES**

FOR USE ONLY IN THE STATES OF TEXAS AND OKLAHOMA

Alligare Dicamba 4 may be applied for weed suppression any time after the sorghum has reached the soft dough stage. An agriculturally approved surfactant may be used to improve performance. For aerial applications use at least 2 gallons of water-based carrier per treated acre.

Do not harvest until 30 days after treatment.

**Broadcast rate per treated acre:** 1/2 pint (1/4 lb. a.i.)

**SMALL GRAINS (WHEAT, BARLEY AND OATS)  
NOT UNDERSEEDED TO LEGUMES**

**IMPORTANT:** Observe all precautions and restrictions. Read and follow cleaning, mixing and application instructions.

**RESTRICTIONS:**

**Pre-Harvest Interval (PHI):** Grain (PHI) 7 days

**If small grains are used for pasture or hay, the following restrictions apply:**

- Animals cannot be removed from treated area for slaughter prior to 30 days after last application.
- There is no waiting period between treatment and grazing for non-lactating dairy animals.
- Treated areas may not be grazed by lactating dairy animals until 7 days after treatment.
- Do not harvest hay from treated areas until 37 days after treatment.

**PRECAUTION:**

- Observe all precautions and restrictions on the labels of products used in tank mix treatments.

**WEEDS CONTROLLED**

Alligare Dicamba 4 or combinations with listed tank mix partners will provide control or suppression of the annual broadleaf weeds listed below. For improved control of listed weeds, it is recommended that Alligare Dicamba 4 be applied in a tank mix with other herbicides. Refer to specific crop for tank mix options.

Alkanet <sup>1</sup>	Knawel (German Moss)	Pigweed, Tumble
Bedstraw, Catchweed <sup>1</sup>	Knotweed, Prostrate	Pineappleweed <sup>1</sup>
Bindweed, Field <sup>2</sup>	Kochia	Plantain, Broadleaf <sup>2</sup>
Buckwheat Tartary	Ladysthumb	Poppy, Red Horned <sup>1</sup>
Buckwheat, Wild	Lambsquarters, Common	Puncturevine <sup>1</sup>
Carpentweed <sup>1</sup>	Lettuce, Miners <sup>1</sup>	Purslane, Common
Chamomile, Corn	Lettuce, Prickly	Radish, Wild <sup>1</sup>
Chervil, Bur <sup>1</sup>	Mallow, Common	Ragweed, Common
Chickweed, Common <sup>1</sup>	Mayweed, Chamomile	Ragweed, Giant
Cockle, Corn	(Dogfennel) <sup>1</sup>	(Buffaloweed) <sup>1</sup>
Cockle, Cow	Mustard, Blue	Rocket, London <sup>1</sup>
Cocklebur, Common	(Purple) <sup>1</sup>	Rocket, Yellow <sup>1</sup>
Cornflower	Mustard, Tansy	Salsify (Goatsbeard) <sup>1</sup>
(Bachelorbutton) <sup>1</sup>	Mustard Treacle <sup>1</sup>	Shepherdspurse <sup>1</sup>
Dandelion, Common <sup>2</sup>	Mustard, Tumble	Smartweed, Green
Dock, Curly <sup>2</sup>	(Jim Hill) <sup>1</sup>	Smartweed, Pennsylvania
Dragonhead, American <sup>1</sup>	Mustard, Wild <sup>1</sup>	Sorrel, Red
Evening Primrose,	Nightshade, Black	(Sheep Sorrel) <sup>1</sup>
Cutleaf <sup>1</sup>	Nightshade, Cutleaf <sup>1</sup>	Sowthistle, Annual
Falseflax, Smallseeded <sup>1</sup>	Nightshade Silverleaf <sup>2</sup>	Starthistle, Yellow <sup>1</sup>
Fiddleneck, (Tarweed) <sup>1</sup>	(White Horsenettle)	Sunflower, Common (Wild)
Flixweed <sup>1</sup>	Pennycress, Field	Thistle, Canada <sup>2</sup>
Fumitory <sup>1</sup>	(Farweed, Frenchweed,	Thistle, Russian
Gromwell, Corn <sup>1</sup>	Stinkweed)	Velvetleaf
Groundsel, Common <sup>1</sup>	Pepperweed, Peppergrass <sup>1</sup>	Vetch <sup>1</sup>
Hempnettle <sup>1</sup>	Pigweed, Redroot	Yarrow, Common <sup>2</sup>
Henbit	(Carelessweed)	
Jacobs Ladder <sup>1</sup>	Pigweed, Rough	

<sup>1</sup> These weeds will be controlled with Alligare Dicamba 4 tank mixtures. Refer to tank mix label for specific weeds controlled.

<sup>2</sup> Alligare Dicamba 4 tank mixes will provide suppression of established broadleaf weeds and control their seedlings.

**RATES AND TIMINGS**

Application of Alligare Dicamba 4 may be made before, during or after planting small grains. For best performance, make applications when weeds are in the 2-3 leaf stage and rosettes are less than 2 inches across. Application of Alligare Dicamba 4 to small grains during periods of rapid growth may result in crop leaning. This condition is temporary and will not reduce crop yields.

Use Alligare Dicamba 4 at 2 to 4 fluid ounces per treated acre in wheat, fall seeded barley, and oats, and at 2 to 3 fluid ounces per treated acre in spring seeded barley. Use the higher level of listed rate ranges when treating difficult to control weeds such as kochia, wild buckwheat, cow cockle, prostrate knotweed, Russian thistle, and prickly lettuce or when dense vegetative growth occurs.

Alligare Dicamba 4 used in a tank mix with other herbicides offers the best spectrum of weed control and herbicide tolerant or resistant weed management. Refer to specific crop for Alligare Dicamba 4 rate and application timing.

For applications prior to the emergence of weeds or when sulfonylurea resistant weeds are present or suspected, use a minimum of 3 fluid ounces per treated acre of Alligare Dicamba 4 with a tank mix herbicide. Non-sulfonylurea herbicides such as 2,4-D or MCPA tank mixed with Alligare Dicamba 4 will offer more consistent control of sulfonylurea resistant weeds.

When tank mixing with sulfonylurea herbicides, such as Ally<sup>®</sup>, Amber<sup>®</sup>, Express<sup>®</sup>, Finesse<sup>®</sup>, Gleam<sup>®</sup> and Harmony<sup>®</sup> Extra, use an agriculturally approved surfactant of at least 80% active ingredient at the rate of 1-4 pints/100 gallons of spray, or not more than 0.25-0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature and difficult to control weeds or dense vegetative growth.

**FALL AND SPRING SEEDED WHEAT**

ALLIGARE DICAMBA 4 MUST BE APPLIED TO FALL SEEDED WHEAT PRIOR TO THE JOINTING STAGE. APPLICATIONS TO SPRING SEEDED WHEAT MUST BE MADE BEFORE WHEAT REACHES THE 6 LEAF STAGE.

**NOTE:** Early developing wheat varieties such as TAM 107, MADISON, or WAKEFIELD must receive application between early tillering and the jointing stage. Care should be taken in staging these varieties to be certain that the application occurs prior to the jointing stage.

**TANK MIX TREATMENTS**

Alligare Dicamba 4 may be tank mixed with one or more of the following herbicides. Read and follow the label of each tank mix product used for precautionary statements, directions for use, weeds controlled, geographic and other restrictions. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

**Broadcast rate per treated acre:**

Apply 2-4 fluid ounces Alligare Dicamba 4 with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D Amine or Ester	2,4-D	4 lb/gal	8-12 fluid oz (0.25-0.375 lb a.i./A) <sup>1</sup>
MCPA Amine or Ester	MCPA	4 lb/gal	8-12 fluid oz (0.25-0.375 lb a.i./A) <sup>1</sup>
Ally <sup>®</sup>	metsulfuron-methyl	60% DF	1/10 oz
Amber <sup>®</sup>	triasulfuron	75% DF	0.28 oz
Express <sup>®</sup>	thifensulfuron + tribenuron-methyl	75% DF	1/6 oz
Finesse <sup>®</sup>	chlorsulfuron + metsulfuron-methyl	75% DF	1/3 oz
Glean <sup>®</sup>	chlorsulfuron	75% DF	1/6 oz
Harmony <sup>®</sup> Extra	thifensulfuron + tribenuron-methyl	75% DF	1/3 oz
bromoxynil	bromoxynil	2 lb/gal	1-1.5 pts
Bronate <sup>®</sup>	bromoxynil + MCPA	4 lb/gal	1-2 pts
Curtail <sup>®</sup>	clopyralid+ 2,4-D	2.38 lb/gal	2-2 2/3pts
clopyralid	clopyralid	3 lb/gal	1/4-1/3 pt
diuron <sup>2</sup>	diuron	80% DF	1/2-1.5 lbs
metribuzin <sup>2</sup>	metribuzin	75% DF	1-10 oz
Dakota <sup>®3</sup>	fenoxaprop-ethyl+MCPA	3.1 lb/gal	16 fluid oz
Tiller <sup>®3</sup>	fenoxaprop-ethyl + MCPA + 2,4D	2.7 lb/gal	1-1.7 pts

<sup>1</sup> When using formulations other than 4 lbs/gal use pounds active/acre listed.

<sup>2</sup> Tank mixtures for fall seeded wheat only.

<sup>3</sup> Use 2 fluid ounces of Alligare Dicamba 4 only. Do not use if wild oats is the target weed. Do not use Alligare Dicamba 4 as a tank mix treatment with Dakota<sup>®</sup> or Tiller<sup>®</sup> on Durum wheat.

**SPECIAL USE TANK MIXES FOR SPRING AND FALL SEEDED WHEAT**

(See Footnotes for Applicable Uses)

Alligare Dicamba 4 may be tank mixed with one or more of the following herbicides. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

**Broadcast rate per treated acre:**

Apply 3-4<sup>1</sup> fluid ounces Alligare Dicamba 4 with:

Product <sup>2</sup>	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D or MCPA Amine	2,4-D or MCPA	4 lb/gal	1-2 pts <sup>3</sup> (0.5-1.0 lb a.i./A) <sup>4</sup>
2,4-D or MCPA Ester	2,4-D or MCPA	4 lb/gal	1-1.5 pts <sup>3</sup> (0.5-.75 lb a.i./A) <sup>4</sup>
Ally <sup>®</sup>	metsulfuron-methyl	60% DF	1/20-1/10 oz
Amber <sup>®</sup>	triasulfuron	75% DF	0.14-0.28 oz
Express <sup>®</sup>	thifensulfuron + tribenuron-methyl	75% DF	1/12-1/6 oz
Finesse <sup>®</sup>	chlorsulfuron + metsulfuron-methyl	75% DF	1/6-1/3 oz
Glean <sup>®</sup>	chlorsulfuron	75% DF	1/6 oz
Harmony <sup>®</sup> Extra	thifensulfuron + tribenuron-methyl	75% DF	1/6-1/3 oz
Metsulfuron-methyl + 2,4-D Amine or Ester <sup>5</sup>	Metsulfuron-methyl + 2,4-D	60% DF + 4 lb/gal	1/20-1/10 oz + 8 fl oz
Amber <sup>®</sup> + 2,4-D Amine or Ester <sup>5</sup>	triasulfuron + 2,4-D	75% DF + 4 lb/gal	0.14-0.28 oz + 8 fl oz
Express <sup>®</sup> + 2,4-D Amine or Ester <sup>5</sup>	(thifensulfuron + tribenuron-methyl)+ 2,4-D	75% DF + 4 lb/gal	1/12-1/6 oz + 8 fl oz
Finesse <sup>®</sup> + 2,4-D Amine or Ester <sup>5</sup>	(chlorsulfuron + metsulfuron-methyl)+2,4-D	75% DF + 4 lb/gal	1/6-1/3 oz + 8 fl oz
Glean <sup>®</sup> + 2,4-D Amine or Ester <sup>5</sup>	chlorsulfuron + 2,4-D	75% DF + 4 lb/gal	1/6 + 8 fl oz
Harmony <sup>®</sup> Extra+2,4-D Amine or Ester <sup>5</sup>	(thifensulfuron + tribenuron-methyl)+ 2,4-D	75% DF + 4 lb/gal	1/6-1/3 oz + 8 fl oz
Glyphosate <sup>6</sup>	glyphosate	3.0 lb/gal	12-16 fl oz

<sup>1</sup> Alligare Dicamba 4 may be used at 6 fluid ounces on fall seeded wheat in Western Oregon as a spring application only. In CO, KS, NM, OK and TX up to 8 fluid ounces of Alligare Dicamba 4 may be applied on fall seeded wheat after it exceeds the 3 leaf stage for suppression of perennial weeds, such as field bindweed. Applications may be made in the fall following a frost but before a killing freeze. Alligare Dicamba 4 may be tank mixed with 2,4-D amine at 8 fluid ounces after wheat begins to tiller. Periods of extended stress such as cold and wet weather may enhance the possibility of crop injury. For fall applications only, do not use if the potential for crop injury is not acceptable.

<sup>2</sup> Do not use low rates of sulfonylurea herbicides, such as Metsulfuron-methyl, Amber, Express, Finesse, Glean, and Harmony Extra on more mature weeds and/or on dense vegetative growth.

<sup>3</sup> NOTE: For use on Fall Seeded Wheat only. Do not use unless potential crop injury will be acceptable.

<sup>4</sup> When using formulations other than 4 lb/gal use pounds active/acre listed.

<sup>5</sup> Use for improved control of Russian thistle, flixweed, gromwell, mayweed and fiddleneck.

<sup>6</sup> Alligare Dicamba 4 may be applied at 2 fluid ounces with any glyphosate formulation labeled for use as a preplant application to small grains with no waiting period prior to planting. Read and follow label directions of the tank mix product for adjuvant use recommendations.

### FALL SEEDED BARLEY

ALLIGARE DICAMBA 4 MUST BE APPLIED TO FALL SEEDED BARLEY PRIOR TO THE JOINTING STAGE.

**NOTE:** For spring barley varieties that are seeded during the winter months or later, follow the rates and timings given for Spring Seeded Barley.

#### TANK MIX TREATMENTS

Alligare Dicamba 4 may be tank mixed with one or more, but not limited to, the following herbicides. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

#### Broadcast rate per treated acre:

Apply 2-4 fluid ounces Alligare Dicamba 4 with:

Product <sup>1</sup>	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D Amine or Ester	2,4-D	4 lb/gal	8 fluid oz (0.25 lb a.i./A) <sup>2</sup>
MCPA Amine or Ester	MCPA	4 lb/gal	8-12 fluid oz (0.25-0.375 lb a.i./A)
metsulfuron-methyl	metsulfuron-methyl	60% DF	1/20-1/10 oz
Amber <sup>®</sup>	triasulfuron	75% DF	0.14-0.28 oz
Express <sup>®</sup>	thifensulfuron + tribenuron-methyl	75% DF	1/12-1/6 oz
Finesse <sup>®</sup>	chlorsulfuron + metsulfuron-methyl	75% DF	1/6-1/3 oz
Glean <sup>®</sup>	chlorsulfuron	75% DF	1/6 oz
Harmony <sup>®</sup> Extra	thifensulfuron + tribenuron-methyl	75% DF	1/6-1/3 oz
metribuzin	metribuzin	75% DF	1-10 oz
bromoxynil	bromoxynil	2 lb/gal	1-1 1/2 pts
Bronate <sup>®</sup>	bromoxynil + MCPA	4 lb/gal	3/4-1 1/2 pts

<sup>1</sup> Do not use low rates of sulfonylureas (metsulfuron-methyl, Amber<sup>®</sup>, Express<sup>®</sup>, Finesse<sup>®</sup>, Glean<sup>®</sup>, and Harmony<sup>®</sup> Extra) on more mature weeds and/or on dense vegetative growth.

<sup>2</sup> When using formulations other than 4 lb/gal use pounds active/acre listed.

### SPRING SEEDED BARLEY

ALLIGARE DICAMBA 4 MUST BE APPLIED BEFORE SPRING SEEDED BARLEY EXCEEDS THE 4 LEAF STAGE.

#### TANK MIX TREATMENTS

Alligare Dicamba 4 may be mixed with one or more of the following herbicides. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

#### Broadcast rate per treated acre:

Apply 2-4 fluid ounces Alligare Dicamba 4 with:

Product <sup>1</sup>	Active Ingredient	Formulation	Amount of Product Per Acre
MCPA Amine or Ester	MCPA	4 lb/gal	8-12 fluid oz (0.25-0.375 lb a.i./A) <sup>2</sup>
metsulfuron-methyl	metsulfuron-methyl	60% DF	1/20-1/10 oz
Amber <sup>®</sup>	triasulfuron	75% DF	0.14-0.28 oz
Express <sup>®</sup>	thifensulfuron + tribenuron-methyl	75% DF	1/12-1/6 oz
Finesse <sup>®</sup>	chlorsulfuron + metsulfuron-methyl	75% DF	1/6-1/3 oz
Glean <sup>®</sup>	chlorsulfuron	75% DF	1/6 oz
Harmony <sup>®</sup> Extra	thifensulfuron + tribenuron-methyl	75% DF	1/6-1/3 oz
metribuzin	metribuzin	75% DF	1-10 oz
bromoxynil	bromoxynil	2 lb/gal	1-1 1/2 pts
Bronate <sup>®</sup>	bromoxynil + MCPA	4 lb/gal	3/4-1 1/2 pts

<sup>1</sup> Do not use low rates of sulfonylureas (metsulfuron-methyl, Amber<sup>®</sup>, Express<sup>®</sup>, Finesse<sup>®</sup>, Glean<sup>®</sup>, and Harmony<sup>®</sup> Extra) on more mature weeds and/or on dense vegetative growth.

<sup>2</sup> When using formulations other than 4 lb/gal use pounds active/acre listed.

### FALL AND SPRING SEEDED OATS

ALLIGARE DICAMBA 4 MUST BE APPLIED BEFORE SPRING SEEDED OATS EXCEED THE 5 LEAF STAGE. APPLICATIONS TO FALL SEEDED OATS MUST BE MADE PRIOR TO THE JOINTING STAGE.

#### TANK MIX TREATMENTS

Alligare Dicamba 4 may be tank mixed with one or more of the following herbicides. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

#### Broadcast rate per treated acre:

Apply 2-4 fluid ounces Alligare Dicamba 4 with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
MCPA Amine or Ester	MCPA	4 lb/gal	8-12 fluid oz (0.25-0.375 lb a.i./A) <sup>1</sup>

<sup>1</sup> When using formulations other than 4 lb/gal use pounds active/acre listed.

### FALL AND SPRING SEEDED TRITICALE EXCEPT CALIFORNIA

#### EARLY SEASON APPLICATIONS

Apply 2-4 fluid ounces of Alligare Dicamba 4 to triticale.

Early season applications to fall-seeded triticale must be made prior to jointing stage.

Early season applications to spring-seeded triticale must be made before triticale reaches the 6-leaf stage.

#### TANK MIXES

Alligare Dicamba 4 may be tank mixed with bromoxynil. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

### SUGARCANE

**IMPORTANT:** Observe all restrictions, precautions, mixing, and application instructions.

Consult your local or state authorities for possible application restrictions, especially concerning aerial applications and advice concerning special local use situations.

Application made over the top of actively growing sugarcane may result in crop injury. When possible, direct the spray beneath the sugarcane canopy in order to minimize the likelihood of crop injury. The use of directed sprays will also aid in maximizing spray coverage of weed foliage.

#### RESTRICTION:

- Do not exceed the maximum single application rate of 2 pints (1.0 lb. a.i.) Alligare Dicamba 4 with no more than 2 applications per year.

#### WEEDS CONTROLLED

Alligare Dicamba 4, when applied at specified rates, will control many ANNUAL, BIENNIAL and PERENNIAL broadleaf weeds commonly found in sugarcane. (Refer to WEED LIST).

#### RATES AND TIMINGS

Application of Alligare Dicamba 4 may be made any time after weeds have emerged and are actively growing but before the close-in stage of sugarcane. Application rates and timing of Alligare Dicamba 4 are given below. Use the higher level of listed rate ranges when treating dense vegetative growth.

Weed Stage & Type	Broadcast Rate Per Treated Acre		PHI
	Amount of Formulated Alligare Dicamba 4 (pints)	Equivalent Lbs. a.i.	
Annual - Small, actively growing - Established weed growth	1/2-1	1/4-1/2	87 days
	1-1 1/2	1/2-3/4	
Biennial	1-2	1/2-1	
Perennial	2	1 <sup>1</sup>	

<sup>1</sup> Application made over the top of actively growing sugarcane may result in crop injury.

#### TANK MIX TREATMENTS

Alligare Dicamba 4 may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

Herbicide	Rate Per Treated Acre (lbs. a.i.)
ametryn	2/5-8
asulam	2-3 1/3
atrazine	2/5-4
2,4-D	1 1/2-3*

\*Application of Alligare Dicamba 4 plus 2,4-D tank mix at the higher listed rate ranges may result in crop injury.

### PASTURE, HAY, RANGELAND, AND FARMSTEAD (Non-Cropland)

**IMPORTANT:** Observe all precautions and restrictions. Read and follow mixing and application instructions.

Alligare Dicamba 4 may be used on pasture, hay, rangeland, farmstead (non-cropland) (including fence rows and non-irrigation ditchbanks) for broadleaf weed and brush control. Alligare Dicamba 4 may also be applied to non-cropland areas for the control of broadleaf weeds in Noxious Weed Control Programs, Districts or Areas including broadcast or spot treatment of roadsides and highways, utilities, railroad and pipeline rights-of-way. Noxious weeds must be recognized at the state level but programs may be administered at state, county or other level.

Alligare Dicamba 4 uses described in this section also pertain to small grains (such as barley, forage sorghum, oats, rye, sudangrass or wheat) grown for pasture use only.

#### PRECAUTIONS:

- NEWLY SEEDED AREAS, including small grains grown for pasture may be severely injured if rates of Alligare Dicamba 4 greater than 1 pint/A are applied.
- ESTABLISHED GRASS CROPS growing under stress can exhibit various injury symptoms that may be more pronounced if herbicides are applied.
- Bentgrass, carpetgrass, buffalograss and St. Augustine grass may be injured at rates exceeding 1 pint Alligare Dicamba 4 (1/2 lb a.i.) per treated acre. Usually colonial bentgrasses are more tolerant than creeping types. Velvetgrasses are most easily injured. Treatments will kill or injure alfalfa, clovers, lespedeza, wild winter peas, vetch and other legumes.

#### RESTRICTIONS:

- ANIMALS CANNOT BE REMOVED FROM TREATED AREA FOR SLAUGHTER PRIOR TO 30 DAYS AFTER LAST APPLICATION.
- THERE IS NO WAITING PERIOD BETWEEN TREATMENT AND GRAZING FOR NON-LACTATING ANIMALS.
- Do not exceed the maximum single application rate of 2 pints (1.0 lb. a.i.) Alligare Dicamba 4 per acre with no more than 2 applications per year.

#### TIMING RESTRICTIONS FOR LACTATING DAIRY ANIMALS FOLLOWING TREATMENT:

Alligare Dicamba 4 Rate Per Treated Acre	Days Before Grazing	Days Before Hay Harvest
Up to 1 pint (1/2 lb. a.i.)	7 days	37 days
Up to 2 pints (1 lb. a.i.)	21 days	51 days

#### MIXING AND APPLICATION

Alligare Dicamba 4 can be applied using water, oil in water emulsions including invert systems, or sprayable fluid fertilizer as a carrier. A COMPATIBILITY TEST (see COMPATIBILITY TEST section) should be made prior to tank mixing.

To prepare oil in water emulsions, half-fill spray tank with water, then add appropriate amount of emulsifier. With continuous agitation, slowly add the herbicide and then the oil (such as diesel oil or fuel oil) or a premix of oil plus additional emulsifier to spray tank. Complete filling of spray tank with water. Maintain vigorous agitation during spray operation to prevent oil and water from forming separate layers.

Alligare Dicamba 4 may be applied broadcast using either ground or aerial application equipment. When using ground equipment, apply 3 to 600 gallons of diluted spray per treated acre. Volume of spray applied will depend on the height, density, and type of weeds or brush being treated and on the type of equipment being used. When using aerial equipment apply 2 to 40 gallons of diluted spray per treated acre in a water-based carrier.

Alligare Dicamba 4 may be applied to individual clumps or small areas (SPOT TREATMENT) of undesirable vegetation using handgun or similar types of application equipment. Apply diluted sprays to allow complete wetting (up to run off) of foliage and stems.

Herbicide adjuvants or other spray additives (emulsifiers, surfactants, wetting agents, drift control agents, or penetrants) may be used for wetting, penetration, or drift control. Spray additives must be agriculturally approved when used in pasture applications. If spray additives are used, read and follow all use recommendations and precautions on product label.

#### WEEDS CONTROLLED

Alligare Dicamba 4, when applied at specified rates, will give control of many ANNUAL, BIENNIAL, and PERENNIAL broadleaf weeds, and many WOODY brush and vine species commonly found in pasture, hay, rangeland, and farmstead (non-cropland) areas. (Refer to WEED LIST). Noted (\*) PERENNIAL weeds may be controlled with lower rates of either Alligare Dicamba 4 or Alligare Dicamba 4 plus 2,4-D. See the following RATES AND TIMINGS section.

**RATES AND TIMINGS**

Application rates and timing of Alligare Dicamba 4 are given below. Use the higher level of listed rate ranges when treating dense or tall vegetative growth.

Weed Stage & Type	Broadcast Rate Per Treated Acre	
	Amount of Formulated Alligare Dicamba 4 (pints)	Equivalent Lbs. a.i.
<b>Annual</b>		
Small, actively growing	1/2-1	1/4-1/2
Established weed growth	1-1 1/2	1/2-3/4
<b>Biennial<sup>1</sup></b>		
Rosette diameter		
Less than 3 inches	1/2-1	1/4-1/2
3 inches or more	2	1
Bolting	2	1
<b>Perennial</b>		
Suppression or top growth control	1-2	1/2-1
Noted (*) Perennials	2	1
Other Perennials	2	1
<b>Woody Brush &amp; Vines</b>		
Top Growth Suppression	1-2	1/2-1
Top Growth Control <sup>2</sup>	2	1
Stems and Stem Suppression	2	1

<sup>1</sup> For best performance, make application when BIENNIAL WEEDS are in the rosette stage.

<sup>2</sup> Species noted in WEED LIST section will require tank mixtures for adequate control.

\* Rates above 1.0 lb a.i./A are spot treatments only.

**TANK MIX TREATMENTS**

Alligare Dicamba 4 may be tank mixed with one or more of the following herbicides for control of grasses, additional broadleaf weeds, and woody brush and vines. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

Herbicide	Rate Per Treated Acre (lbs. a.i.)
Pasture, hay, rangeland and farmstead (non-cropland) use:	
glyphosate	3/4 - 3 3/4
metsulfuron methyl	0.0038-0.011
paraquat	1/2 - 1
picloram	1/8 - 3
triclopyr	3/4 - 9
2,4-D	1/4 - 6

Due to the variations that may occur in formulated products and specific use ingredients (e.g. water supplies), a COMPATIBILITY TEST is recommended prior to actual tank mixing.

**CUT SURFACE TREE TREATMENTS**

Alligare Dicamba 4 may be applied as a cut surface treatment for control of unwanted trees and prevention of sprouts of cut trees. A mix of 1 part Alligare Dicamba 4 with 1 to 3 parts water should be used in application. Use the lower dilution when treating difficult-to-control species.

**FRILL OR GIRDLING TREATMENTS:** Make a continuous cut or a series of overlapping cuts using an axe to girdle tree trunk. Spray or paint cut surface with the Alligare Dicamba 4/water mix.

**STUMP TREATMENTS:** Spray or paint freshly cut surface with the water mix. The area adjacent to the bark should be thoroughly wet.

**Note:** For more rapid foliar effects, 2,4-D may be added to the Alligare Dicamba 4/water mix.

**DORMANT APPLICATIONS FOR CONTROL OF MULTIFLORA ROSE**

Alligare Dicamba 4 can be applied when plants are dormant as an undiluted SPOT-CONCENTRATE directly to the soil or as a LO-OIL BASAL BARK treatment using an oil-water emulsion solution.

SPOT-CONCENTRATE applications of Alligare Dicamba 4 should be applied directly to the soil as close as possible to the root crown but within 6-8 inches of the crown. On sloping terrain, application should be made to the uphill side of the crown. Do not make application when snow or water prevents applying Alligare Dicamba 4 directly to the soil. The use rate of Alligare Dicamba 4 is dependent on the canopy diameter of the multiflora rose. Examples: Use Alligare Dicamba 4 at 1/4, 1 or 2 1/4 fluid ounces of product respectively, for 5, 10 or 15 feet canopy diameters. Do not exceed a total of 2 quarts Alligare Dicamba 4 per acre per year.

LO-OIL BASAL BARK applications of Alligare Dicamba 4 should be applied to the basal stem region from the ground line up to a height of 12 to 18 inches. Spray until runoff, with special emphasis on covering the root crown. For best results, make application when plants are dormant. Do not make application after bud break or when plants are showing signs of active growth. Do not make application when snow or water prevents applying Alligare Dicamba 4 to the ground line. Refer to Mixing and Applications above in this section for method of preparing oil-in-water emulsion. Example for making approximately 2 gallons of a LO-OIL spray solution mixture: combine 1 1/2 gallons water plus 1 ounce emulsifier plus 1 pint Alligare Dicamba 4 plus 2 1/2 pints of No. 2 diesel fuel. Adjust amounts of materials used proportionately to the amount of final spray solution desired. Do not exceed 8 gallons of spray solution mix applied per acre per year.

**CONSERVATION RESERVE PROGRAM (CRP) ACRES**

**IMPORTANT:** Observe all precautions, restrictions, mixing and application directions.

Alligare Dicamba 4 can be used on both newly seeded and established grasses grown in Conservation Reserve or Federal Set-Aside Programs. For program lands, such as Conservation Reserve Program, consult program rules to determine whether grass or hay may be used. The more restrictive requirements of the program rules or this label must be followed.

Agriculturally approved surfactants may be added to the spray mixture to improve postemergence weed control, particularly in dry growing conditions.

**NEWLY SEEDING AREAS**

Alligare Dicamba 4 may be applied either preplant or postemergence to newly seeded grasses or small grains such as barley, oats, rye, sudangrass, wheat, or other grain species grown as a cover crop. Postemergence applications may be made after seedling grasses exceed the 3-leaf stage. Rates of Alligare Dicamba 4 greater than 1 pint per treated acre may severely injure newly seeded grasses. Preplant applications - injury to new seedlings may occur if intervals between application and grass planting is less than 45 days per pint of Alligare Dicamba 4 per treated acre West of the Mississippi River or 20 days per pint East of the Mississippi River.

**ESTABLISHED GRASS STANDS**

Established grass stands are perennial grasses planted one or more seasons prior to treatment. Certain species: bentgrass, carpetgrass, smooth brome, buffalograss or St. Augustine grass may be injured when treated with Alligare Dicamba 4 at rates exceeding 1 pint per treated acre.

**WEEDS CONTROLLED**

Alligare Dicamba 4, when applied at specified rates, will control many annual and biennial weeds and provide control or suppression of many perennial weeds. (Refer to WEED LIST).

**RATES AND TIMINGS**

Application rates and timing of Alligare Dicamba 4 treatment are given below. Use the higher rate of the rate range when vegetation is either dense or tall, or when weeds are growing under stressed conditions such as drought or cool temperature.

**RESTRICTIONS:**

- Do not exceed the maximum single application rate of 2 pints (1.0 lb. a.i.) Alligare Dicamba 4 per acre with no more than 2 applications per year.
- Do not use adjuvants containing penetrants such as petroleum based oils after grass emergence on newly seeded grasses.

**PRECAUTION:**

- Alligare Dicamba 4 treatment will injure or may kill alfalfa, clovers, lespedeza, wild winter peas, vetch, and other legumes.

Weed Stage & Type	Broadcast Rate Per Treated Acre		PHI
	Amount of Formulated Alligare Dicamba 4 (pints)	Equivalent lbs. a.i.	
<b>Annual</b>			
Small, actively growing	1/4-1	1/8-1/2	For grass forage: 0 days
Established weed growth	1	1/2	
<b>Biennial<sup>1,2</sup></b>			For grass hay: 7 days
Rosette diameter			
Less than 3 inches	1/2-1	1/4-1/2	
3 inches or greater	1-2	1/2-1	
Bolting biennial	2	1	
<b>Perennial<sup>2</sup></b>			
Suppression/Control	2	1	

<sup>1</sup> For best results, treat Biennial weeds with Alligare Dicamba 4 when they are in the rosette stage of growth.

<sup>2</sup> Biennial and Perennial weeds will require follow-up (sequential) treatments for seedling control and escapes.

**TANK MIX TREATMENTS**

To control grasses and additional broadleaf weeds, Alligare Dicamba 4 may be tank mixed with other herbicides registered for use in Conservation Reserve Programs such as 2,4-D, glyphosate, paraquat, metsulfuron, and others. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

**ASPARAGUS**

FOR USE ONLY IN THE STATES OF CALIFORNIA, OREGON, AND WASHINGTON

**IMPORTANT:** Observe all restrictions, precautions, mixing, and application instructions.

**NOTE:** If spray contacts emerged spears, crooking (twisting) of some spears may result. If such crooking occurs, discard affected spears.

**RESTRICTIONS:**

- Do not harvest prior to 24 hours after treatment.
- Do not use in the Coachella Valley of California.
- A retreatment may be made if needed; however, DO NOT EXCEED a total of 1 pint (1/2 lb. a.i.) of Alligare Dicamba 4 per treated acre per crop year.

**RATES AND TIMINGS**

Apply Alligare Dicamba 4 to emerged and actively growing weeds in 40 to 60 gallons of diluted spray per treated acre immediately after cutting the field, but at least 24 hours before the next cutting.

**TANK MIXING**

Alligare Dicamba 4 may be applied in a tank mixture with either 2,4-D or glyphosate herbicide for improved control of noted (\*) weeds. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

Weeds	Rate Per Treated Acre
Mustard, Black	1/2-1 pt. (1/4-1/2 lb. a.i.)
Pigweed, Redroot (Carelessweed)	
Sowthistle, Annual	
*Thistle, Canada	
*Thistle, Russian	
*Bindweed, Field	1 pt. (1/2 lb. a.i.)
Chickweed, Common	
Goosefoot, Nettleleaf	
Radish, Wild	
Thistle, Milk	

**TURF AND LAWN**  
FOR USE IN FARMSTEAD (NON-CROPLAND) AND SOD FARMS

**IMPORTANT:** Observe all restrictions, precautions, mixing, and application instructions.

To avoid injury to newly seeded grasses, application of Alligare Dicamba 4 should be delayed until after the second mowing. Further-more, application rates in excess of 1 pint (1/2 lb. a.i.) per treated acre may cause noticeable stunting or discoloration of sensitive grass species such as bentgrass, carpetgrass, buffalograss, and St. Augustine grass.

In areas where roots of sensitive plants extend, do not apply in excess of 1/4 pint (1/8 lb. a.i.) of Alligare Dicamba 4 per treated acre on coarse textured (sandy-type) soils, or in excess of 1/2 pint (1/4 lb. a.i.) per treated acre on fine textured (clayey-type) soils. Do not make repeat applications in these areas for 30 days and until previous applications of Alligare Dicamba 4 have been activated in the soil by rain or irrigation.

**WEEDS CONTROLLED**

Alligare Dicamba 4, when applied at specified rates, will give control of many ANNUAL, BIENNIAL, and noted (\*) PERENNIAL broadleaf weeds commonly found in turf. Alligare Dicamba 4 will also give growth suppression of many other listed PERENNIAL broadleaf weeds and WOODY brush and vine species. (Refer to WEED LIST).

**MIXING AND APPLICATION**

Apply 30 to 200 gallons of diluted spray per treated acre (3 qts. to 4 1/4 gals. per 1,000 sq. ft.), depending on density or height of weeds treated and on the type of equipment used.

**RATES AND TIMINGS**

Use the higher level of listed rate ranges when treating dense vegetative growth. For best performance, apply when weeds are emerged and actively growing.

**RESTRICTION:**

- Do not exceed the maximum single application rate of 2 pints (1.0 lb. a.i.) of Alligare Dicamba 4 per acre with no more than 2 applications per year.

Weed Stage & Type	Alligare Dicamba 4		
	Pints per treated acre	Lbs. a.i. per treated acre	Teaspoons per 1,000 sq. ft.
<b>Annual</b>			
Small, actively growing	1/4-1	1/4-1/2	1-2 1/4
Established weed growth	1-1 1/2	1/2-3/4	2 1/4-3 1/4
<b>Biennial Rosette diameter</b>			
Less than 3 inches	1/2-1	1/4-1/2	1-2 1/4
3 inches or more	1-2	1/2-1	2 1/4-4 1/2
<b>Perennial and Woody</b>			
Brush and Vines	1-2	1/2-1	2 1/4-4 1/2

**TANK MIX TREATMENTS**

Alligare Dicamba 4 may be tank mixed with 2,4-D, MCPA, MCPP, or bromoxynil for control of additional weeds listed on the tank mix product label. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

Apply 1/5 to 1/2 pint (1/10 to 1/4 lb. a.i.) of Alligare Dicamba 4 per treated acre with the labeled rate of 2,4-D, MCPA, MCPP, or bromoxynil.

**RESTRICTION:**

- Repeat treatments may be made as needed; however, DO NOT EXCEED 2 pints (1 lb. a.i.) of Alligare Dicamba 4 per treated acre during the growing season.

**GRASS SEED CROPS**

GRASSES GROWN FOR SEED SUCH AS BERMUDA GRASS, BLUEGRASS, FESCUE AND RYEGRASS

**IMPORTANT:** Observe all restrictions, precautions, mixing, and application instructions.

Refer to the PASTURE, HAY, RANGELAND, AND FARMSTEAD (NONCROPLAND AREAS) section for possible grazing and feeding restrictions.

**WEEDS CONTROLLED**

Alligare Dicamba 4 will provide control or suppression of annual broadleaf weeds listed below. For improved control of listed weeds plus additional weeds, it is recommended that Alligare Dicamba 4 be applied in a tank mix with other herbicides.

Alfalfa <sup>1</sup>	Clover	Ladythumb
Bedstraw, Catchweed	Cockle, White	Lambsquarters, Common
Bindweed, Field	Dock, Broadleaf	Lettuce, Prickly
Buttercup, Corn	Dock, Curly	Mayweed (Dogfennel)
Buttercup, Creeping	Hemlock, Poison	Ragwort, Tansy
Buttercup, Western Field	Knawelweed, Russian <sup>1</sup>	Sorrel, Red (Sheep Sorrel)
Catchfly, Nightflowering	Knawel	Sowthistle, Annual
Chamomile, Corn	Kochia	Starwort, Little
Chickweed, Common	Knotweed, Prostrate	Thistle, Canada <sup>1</sup>
Chickweed, Mouseear		

<sup>1</sup> Top growth control only

**RATES AND TIMINGS**

Apply 1/2 to 1 pint of Alligare Dicamba 4 per treated acre on SEEDLING GRASS after the crop reaches the 3-5 leaf stage.

Apply up to 2 pints of Alligare Dicamba 4 on well-established perennial grass. DO NOT APPLY AFTER THE GRASS SEED CROP BEGINS TO JOINT. For best performance, make applications when weeds are in the 2-4 leaf stage and rosettes are less than 2 inches across. Use the higher level of listed rate ranges when treating more mature weeds or dense vegetative growth.

**RESTRICTIONS:**

- Do not exceed the maximum single application rate of 2 pints (1.0 lb. a.i.) of Alligare Dicamba 4 per acre with no more than 2 applications per year.
- Do not use on bentgrass unless possible crop injury can be tolerated.

**TANK MIX TREATMENTS**

For control of grasses or additional broadleaf weeds, Alligare Dicamba 4 may be tank mixed with all broadleaf herbicides registered for use in Grass Seed Production. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

**Broadcast Rate Per Treated Acre:**

Apply 1/2 to 2 pints Alligare Dicamba 4 with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
2,4-D Amine or Ester	2,4-D	4 lb/gal	1-4 pts. (0.5-2.0 lb a.i./A) <sup>1</sup>
MCPA Amine	MCPA	4 lb/gal	1-2 pts (0.5-1.0 lb a.i./A) <sup>1</sup>
bromoxynil	bromoxynil	2 lb/gal	1-2 pts
Curtail®	clopyralid + 2,4-D	2.38 lb/gal	1 3/4-4 pts
diuron	diuron	80% DF	2-4 lbs
clopyralid	clopyralid	3 lb/gal	1/4-1 pt

<sup>1</sup> When using formulations other than 4 lb/gal use pounds active/acre listed.

**ANNUAL GRASS CONTROL**

For suppression of ANNUAL GRASS WEEDS such as:

- Brome, Downy (Cheatgrass)
- Brome, Ripgut
- Fescue, Rattail
- Windgrass

Apply up to 2 pints (1lb. a.i.) of Alligare Dicamba 4 per treated acre in the fall or late summer after harvest and burning of established grass seed crops (maximum of 2 treatments per year). Applications should be made immediately following first irrigation when the soil is moist and before weeds have more than 2 leaves.

**PREPLANT DIRECTIONS (POST HARVEST/FALLOW/CROP STUBBLE/SET-A-SIDE)  
FOR BROADLEAF WEED CONTROL BEFORE WHEAT, CORN, SORGHUM, SOYBEANS**

**IMPORTANT:** Observe all restrictions, precautions, mixing, and application instructions.

**WEEDS CONTROLLED**

Alligare Dicamba 4 may be applied alone or in tank mix combinations with other herbicides registered for this use.

Alligare Dicamba 4 can be applied either POST HARVEST in the fall, spring or summer during the FALLOW period or to CROP STUBBLE/SET-A-SIDE acres. Alligare Dicamba 4, when applied at the specified rates, will control many ANNUAL broadleaf weeds; see the WEEDS CONTROLLED section under small grains. In addition, Alligare Dicamba 4 will control or suppress the following BIENNIAL and PERENNIAL broadleaf weeds:

Alfalfa <sup>1</sup>	Dock, Curly <sup>1</sup>	Sowthistle, perennial <sup>1</sup>
Artichoke, Jerusalem	Dogbane, Hemp	Spurge, leafy
Bindweed, Field	Garlic, Wild <sup>2</sup>	Thistle Bull
Bindweed, Hedge	Horsenettle, Carolina	Thistle, Canada <sup>2</sup>
Blueweed, Texas	Knawelweed, Diffuse	Thistle, Milk
Bursage	Knawelweed, Spotted	Thistle, Musk
(Bur Ragweed)	Nightshade, Silverleaf	Thistle, Plumeless
(Povertyweed)	Redvine	Thistle, Scotch
(Lakewood) <sup>1</sup>	Smartweed, Swamp	Trumpet creeper (Buckvine)
Dandelion, Common <sup>1</sup>		

<sup>1</sup> Perennials may be controlled using Alligare Dicamba 4 at rates lower than those listed for other listed perennial weeds. (See RATES AND TIMINGS under this heading.)

<sup>2</sup> See the SPECIAL TANK MIX TREATMENTS section under this heading for specific control programs for these weeds.

**RATES AND TIMINGS**

Apply Alligare Dicamba 4 as a broadcast or spot treatment to emerged and actively growing weeds after crop harvest (post harvest) and before a killing frost or in the fallow cropland or crop stubble the following spring or summer. Agriculturally approved spray additives, such as surfactants or oils, may be used to enhance spray coverage and the herbicide's penetration of weed foliage. See Cropping restrictions for recommended interval between application and planting to prevent crop injury.

For best performance, make application when ANNUAL weeds are less than 6 inches tall, when BIENNIAL weeds are in the rosette stage and to PERENNIAL weed regrowth in late summer or fall following a mowing or tillage treatment. Most effective control of upright perennial broadleaf weeds, such as Canada thistle and Jerusalem artichoke, occurs if application is made when the majority of weeds, such as field bindweed and hedge bindweed, are best controlled when weeds are in or beyond the full bloom stage.

Avoid disturbing treated areas following application. Treatments may not kill weeds which develop from seed or underground plant parts, such as rhizomes or bulbets, after the effective period for Alligare Dicamba 4. For seedling control, a follow-up program or other cultural practices could be instituted. For small grain in-crop uses of Alligare Dicamba 4, see the RATE AND TIMINGS section under the SMALL GRAINS heading for details.

**ALLIGARE DICAMBA 4 RATES PER TREATED ACRE**

**RESTRICTION:**

- Do not exceed the maximum single application rate of 2 pints (1.0 lb. a.i.) of Alligare Dicamba 4 per acre with no more than 2 applications per year.

WEED TYPE	AMOUNT OF PRODUCT PER ACRE PER APPLICATION
Annual	1/2-1 pt (8-16 fl. oz.)
Biennial	1-2 pts (16-32 fl. oz.)
Perennial	1-4* pts (16-64 fl. oz.)
Perennial suppression	1-2 pts (16-32 fl. oz.)
Noted (1) perennials	2 pts (32 fl. oz.)
Other perennials	2 pts (32 fl. oz.)

**TANK MIX TREATMENTS**

Alligare Dicamba 4 may be tank mixed with one or more of the following herbicides for control of grasses or additional broadleaf weeds. It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).



**ALLIGARE DICAMBA 4 BROADCAST RATE PER TREATED ACRE FOR ANNUAL WEED CONTROL:**

Apply 1/4 to 1 pint Alligare Dicamba 4 with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
Atrazine <sup>1</sup>	atrazine	4 lb/gal	1/2-6 pts
		90% DF	1/2-3.3 lbs
metsulfuron-methyl <sup>2</sup>	metsulfuron-methyl	75% DF	0.1 oz
Amber <sup>® 2</sup>	triasulfuron	75% DF	0.28-0.35 oz
		paraquat	1-2 pts
paraquat	paraquat	2.5 lb/gal	1.5 pts
		75% DF	0.2 oz
Finesse <sup>® 2</sup>	chlorsulfuron + metsulfuron-methyl		
pronamide <sup>1</sup>	pronamide	50-W	1/2-1.0 lb
Fallow Master <sup>®</sup>	glyphosate + dicamba	1.6 lb/gal	22-44 fluid oz
Landmaster <sup>®</sup> BW	glyphosate + 2,4-D	2.4 lb/gal	27-54 fluid oz
glyphosate	glyphosate	3 lb/gal	8-48 fluid oz
metribuzin <sup>1</sup>	metribuzin	75% DF	1/2-1 lb
		4 lb/gal	3/4-1 1/2 pts
2,4-D	2,4-D	4 lb/gal	1-2 pts (0.5-1 lb a.i./A) <sup>3</sup>

<sup>1</sup> Tank mixes of Alligare Dicamba 4 with these products may be subject to special restrictions. See the Product Label of the tank mix partner for intended use rates, restrictions and other precautions.

<sup>2</sup> When tank mixing with sulfonyleurea herbicides, refer to the product label for rates and restrictions. Use a surfactant of at least 80% active ingredient at the rate of 1-2 quarts/100 gallons of spray or not more than 0.25-0.5% by volume. Use the highest rate of surfactant when using the lower rate ranges of the tank mix and/or when treating more mature weeds or dense vegetative growth. Sulfonyleurea resistant weeds may not be controlled by tank mixes of Alligare Dicamba 4 and a sulfonyleurea. Refer to the Alligare Dicamba 4 tank mix section for alternative tank mixes.

<sup>3</sup> When using formulations other than 4 lb/gal use pounds active/acre listed.

**ALLIGARE DICAMBA 4 BROADCAST RATE PER TREATED ACRE FOR BIENNIAL AND PERENNIAL WEED CONTROL:**

Apply 1 to 2 pints (0.5-1.0 lb. a.i.) of Alligare Dicamba 4 with:

Product	Active Ingredient	Formulation	Amount of Product Per Acre
Curtail <sup>®</sup>	clopyralid + 2,4-D	2.38 lb/gal	2-4 pts
2,4-D	2,4-D	4 lb/gal	2-6 pts (1.0-3 lb a.i./A) <sup>1</sup>
Landmaster <sup>®</sup> BW	glyphosate + 2,4-D	2.4 lb/gal	54 fluid oz
glyphosate	glyphosate	3.0 lb/gal	1-5 qts
picloram	picloram	2 lb/gal	1/2-1 pt

<sup>1</sup> When using formulation other than 4 lb/gal use pounds active/acre listed.

**SPECIAL TANK MIX TREATMENTS**

For suppression of perennial weeds, apply 1/2-1 pint of Alligare Dicamba 4 with 8-16 fluid ounces of glyphosate herbicide per treated acre.

For wild garlic control, apply 1 pint Alligare Dicamba 4 with 3 pints of 2,4-D LV Ester (4 lb/gal) per treated acre. Apply when wild garlic is 4 to 8 inches tall.

For Canada thistle control, use Alligare Dicamba 4, or Alligare Dicamba 4 plus Curtail<sup>®</sup> or Alligare Dicamba 4 plus glyphosate herbicide or glyphosate tank mix treatments.

Application may be made during fallow periods for control of volunteer barley, bulbous bluegrass, downy brome, jointed goatgrass, common rye and volunteer wheat when they are actively growing. Use 1 pint Alligare Dicamba 4 with 1/2-1 lb pronamide 50W. Fall seeded wheat may be planted 9 months or more after application. For best performance, make application between mid-October and mid-December, prior to soil freeze up.

During fallow periods, apply Alligare Dicamba 4 plus Landmaster<sup>®</sup> BW or Fallow Master<sup>®</sup> herbicide to give improved control of kochia, wild buckwheat, prickly lettuce, field bindweed and Canada thistle. Use 1/8-1/4 pint of Alligare Dicamba 4 plus 22 to 54 fluid ounces of Landmaster<sup>®</sup> BW or Fallow Master<sup>®</sup> herbicide for annual weed control or 1/4 to 1/2 pint Alligare Dicamba 4 plus 22 to 54 fluid ounces of Landmaster<sup>®</sup> BW or Fallow Master<sup>®</sup> herbicide for perennial weed suppression.

**CROPPING RESTRICTIONS**

The following use directions are based on a maximum single application rate of 1.0 lb. ae per acre and a maximum annual rate of 2.0 lbs. ae per acre per year.

CORN, SORGHUM and SOYBEANS may be planted in the spring following applications made during the previous year. If less than 1 inch of rainfall occurs between application and first killing frost, cultivate treated areas to allow herbicide to come in contact with moist soil. Cultivation may take place before or immediately after ground thaw.

Soybean injury may occur if the interval between application and planting is less than specified. In areas with greater than 30 inches of rainfall, delay planting for 30 days per pint of Alligare Dicamba 4 per treated acre. In areas with less than 30 inches of rainfall, delay planting for 45 days per pint of Alligare Dicamba 4 per treated acre. Exclude days when ground is frozen.

WHEAT may be planted in the fall or spring following applications. Also, spot applications may be made any time prior to crop emergence if crop injury can be tolerated in treated areas. Wheat injury may occur if the interval between application and planting is less than specified.

East of the Mississippi River, the interval is 20 days per pint of Alligare Dicamba 4 per treated acre or 1.25 days per 1 ounce. Moisture is essential for Alligare Dicamba 4 degradation. Exclude days when ground is frozen.

West of the Mississippi River, the interval is 45 days per pint of Alligare Dicamba 4 per treated acre or 3 days per ounce. Moisture is essential for Alligare Dicamba 4 degradation. Exclude days when ground is frozen.

Following a normal harvest of barley, oats, or wheat, any rotation crop may be planted. If the interval before harvest is shortened, such as when cover crops will be plowed under, do not follow up with the planting of a sensitive crop (beans, cotton, flowers, fruit trees, grapes, ornamentals, peas, potatoes, soybeans, sunflowers, tobacco and tomatoes).

**CONTROL OF PERENNIAL BROADLEAF WEEDS IN CROPLAND (SPOT APPLICATION ONLY)**

FOR USE ONLY IN THE STATES OF IDAHO, MONTANA, NEVADA, OREGON, UTAH AND WASHINGTON.

**IMPORTANT:** Observe all restrictions, precautions, mixing, and application instructions.

**RESTRICTIONS:**

- Do not treat sub-irrigated cropland or areas where the soil remains saturated with water throughout the year.
- Do not exceed the maximum single application rate of 2 pints (1.0 lb. a.i.) of Alligare Dicamba 4 per acre with no more than 2 applications per year.

**NOTE:** Do not use unless injury to wheat or rotated barley will be acceptable. Application may be made up to one month prior to planting of wheat.

**WEEDS CONTROLLED**

Alligare Dicamba 4, when applied at specified rates, will control many broadleaf weeds including:

Bindweed, Field	Knapweed, Russian
Dock, Broadleaf (Bitterdock)	Ragwort, Tansy
Dock, Curly	Spurge, Leafy
Knapweed, Black	Thistle, Canada

**RATES AND TIMINGS**

Alligare Dicamba 4 may be applied at any time following a crop harvest to stubble, fallow or other cropland. Application should be made when weeds are actively growing and prior to a killing frost.

Barley, oats, corn, sorghum (milo), annual or perennial grass crops may be planted into treated areas one year after application. Crops grown for seed (other than perennial grass seed) should not be planted into treated areas until three years after application. Do not plant broadleaf crops such as alfalfa, beans, peas, potatoes, or sugarbeets into treated areas until two years after application.

In most cases, treatments will not kill perennial weed seedlings, which germinate from seed one or two years after treatment. Once the effect of the chemical has been lost, a follow-up program for seedling control or other cultural practices should be instituted.

**WIPER APPLICATION USES**

**IMPORTANT:** Observe all restrictions, precautions, mixing, and application instructions.

Alligare Dicamba 4 may be applied through wiper application equipment to control or suppress actively growing broadleaf weeds, brush and vines. Use a solution containing 1 part Alligare Dicamba 4 to 1 part water. Do not contact desirable vegetation with herbicide solution. Make wiper application only to crops (including pastures) and non-cropland areas described in this label with the exception of Grain Sorghum (Milo).

**RIGHTS-OF-WAY, UTILITY AND INDUSTRIAL AREAS, AND FENCEROWS**

Alligare Dicamba 4 may be used for use on non-crop land areas such as rights-of-way (such as roadways, rest areas, utility, railroad, highway, pipeline, and rights-of-way that run through pasture and rangeland), utility facilities (such as substations, pipelines, tankfarms, pumping stations, parking and storage areas, fencerows, and non-irrigated ditchbanks), brush control for forest site preparation or maintenance.

Observe all Precautions on this label. Read and follow the Mixing and Application section.

**RESTRICTION:**

- Retreatments may be made as needed, however, do not exceed a total of 4 pts (2 lbs a.i.) of Alligare Dicamba 4 per treated acre during a growing season.

**PRECAUTION:**

- Herbicide adjuvants or other spray additives (emulsifiers, spreader stickers, surfactants, wetting agents, drift control agents, or penetrants) may be used for wetting, penetration, or drift control. Spray additives must be agriculturally approved when used in pasture applications. If spray additives are used, read and follow all use recommendations and precautions on product label.

**Rights-of-Way**

Alligare Dicamba 4 can be used to control many broadleaf weeds on rights-of-way. This use includes applications to roadside, roadway and highways, to areas along utilities such as cable and powerlines, railroad track and embankment, highways, highway medians, bridge abutments, pipelines, and rights-of-way that run through pasture and rangeland. Use controlled application techniques that minimize the risk of off-target movement.

**Utility and Industrial Areas**

Alligare Dicamba 4 can be used to control many broadleaf weeds and brush in noncrop areas on or surrounding substations, pipelines, tankfarms, pump stations, production facilities, and bareground situations. It may also be used on parking and storage areas (refer to Best Stewardship Practices to avoid direct runoff from impervious surfaces).

**Fencerows**

Alligare Dicamba 4 can be used to control many broadleaf weeds and brush in fencerows.

**Mixing and Application**

Alligare Dicamba 4 can be applied using water, oil in water emulsions including invert systems, or sprayable fluid fertilizer as a carrier. A compatibility test (see Compatibility Test section) should be made prior to tank mixing.

To prepare oil in water emulsions, half-fill spray tank with water, then add the appropriate amount of emulsifier. With continuous agitation, slowly add the herbicide and then the herbicidal oil or a pre-mix of oil plus additional emulsifier to spray tank. Complete filling of spray tank with water. Maintain vigorous agitation during spray operation to prevent oil and water from forming separate layers.

Alligare Dicamba 4 may be applied broadcast using either ground or aerial application equipment. When using ground equipment, apply low or high volume sprays of between 3-600 gals, of diluted spray per treated acre. Volume of spray applied will depend on the height, density, and type of weeds or brush being treated and on the type of equipment being used. When using aerial equipment, apply 5-40 gals, of diluted spray per treated acre.

Alligare Dicamba 4 may be applied to individual clumps or small areas (spot treatment) of undesirable vegetation using handgun or similar types of application equipment. Apply diluted spray to allow complete wetting (up to runoff) of foliage and stems.

**Weeds and Brush Controlled**

Alligare Dicamba 4, when applied at specified rates, will give control of many annual, biennial, and perennial broadleaf weeds, and many woody brush and vine species commonly found in non-crop land areas (Refer to Weed List). Noted (\*) perennial weeds may be controlled with lower rates of either Alligare Dicamba 4 or Alligare Dicamba 4 plus tank mix combinations. See Rates and Timings below.

**Rates and Timings**

Application rates and timings of Alligare Dicamba 4 are given below. Use the higher level of listed rate ranges when treating dense or tall vegetative growth.

Weed Stage and Type	Amount of Product Per Acre	Gals of Spray Mixture Per Acre**	Spray Concentration for Low Volume Application**** (% vol/vol)
Annual Small, Actively Growing Established weed growth	1/2 - 1 pt	25 - 50	3
	1 - 1 1/2 pts	50 - 75	3
Biennial* - Rosette diameter Less than 3" 3" or more Bolting	1/2 - 1 pt	25 - 50	3 - 4
	1 - 2 pts	50 - 100	3 - 4
	2 - 3 pts	100 - 150	3 - 4
Perennial Suppression or top growth control Noted (*) Perennials	1/2 - 1 pt	50 - 100	4
	2 - 4 pts	100 - 200	4
	4 pts	200	5
Woody Brush and Vines*** Top Growth Stems And Roots	1/2 - 4 pts	50 - 200	5
	4 pts	200	5

\* For best performance, make application when biennial weeds are in the rosette stage.

\*\*Assuming typical application rate of 1 qt of Alligare Dicamba 4/100 gals.

\*\*\*Tank mixes may be required for optimal control. Refer to Weed List.

\*\*\*\*Low volume rates must not exceed 4 pts of Alligare Dicamba 4 maximum per acre per year (5% volume/volume = 10 gals, maximum solution per acre per year).

**Tank Mix Options for Rights-Of-Way, Utility, and Industrial Areas, and Fencerows**

Alligare Dicamba 4 may be tank mixed with other herbicides for additional weed control. The following table lists example options, but does not limit tank mix options.

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions and precautionary language of the products in the mixture (for example, first aid from one product, spray drift management from another).

Herbicide	Rates Per Treated Acre (lbs a.i.)
norflurazon (Predict®)	Consult product labels for rate recommendations.
proflaminate (Endurance®, Alligare Proflaminate 65 WG Herbicide, Alligare Proflaminate 4L)	
glufosinate (Finale®)	
glyphosate (Roundup®, Accord®, Glyphosate 5.4, Glyphosate 4 Plus, Alligare Glyphosate 4)	
metsulfuron methyl (Escort®, Alligare MSM 60)	
pendimethalin (Pendulum®)	
triclopyr (Redeem®, Garlon®, Alligare Triclopyr 4, Alligare Triclopyr 3, Boulder)	
clopyralid (Transline®, Alligare Clopyralid 3)	
bromacil (Hyvar®, Bromacil 80)	
chlorsulfuron (Telar®, Alligare Chlorsulfuron 75)	
diquat (Reward®, Alligare Diquat Herbicide)	
simazine (Princep®)	
diuron (Karmex®, Alligare Diuron 80DF, Alligare Diuron 4L)	
fosamine ammonium (Krenite®)	
hexazinone (Velpar®, Alligare Hexazinone 75 ULW Herbicide)	
imazapyr (Arsenal®, Alligare Rotary 2 SL, Alligare Ecomazapyr 2 SL, Alligare Imazapyr 2 SL, Alligare Imazapyr 4 SL)	
Imazapic-ammonium (Plateau®, Panoramic 2SL)	
sulfometuron methyl (Oust®, Alligare SFM 75)	
sulfosate (Touchdown®)	
tebuthiuron (Spike®, Alligare Tebuthiuron 80 WG, Alligare Tebuthiuron 20 P)	
2,4-D	

Due to the differences that may occur between specific formulated products and specific use ingredients (e.g., water supplies), a compatibility test (see Compatibility Test section) is recommended prior to actual tank mixing.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

### PESTICIDE STORAGE

Store in original containers in a well-ventilated area separately from fertilizer, feed and foodstuffs. Avoid cross-contamination with other pesticides. Spillage or leakage should be contained and absorbed with clay granules, sawdust, or equivalent material for disposal.

### PESTICIDE DISPOSAL

Triple rinse pesticide from containers and use rinsates in the pesticide application. Wastes which cannot be used according to label instructions may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL Non-refillable container. Plastic or Metal.** Do not reuse or refill this container. Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities, such as burning of plastic containers. If burned, stay out of smoke. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. Triple rinse as follows:

**Non-refillable container less than or equal to 5 gallons:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into formulation equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

**Non-refillable container greater than 5 gallons:** Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution for 30 seconds. Stand the container on its end and tip back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into formulation equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

**Pressure rinse as follows (all sizes):** Empty the remaining contents into formulation equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over formulation equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable container:** Refill this container with pesticide only. Do not reuse this container for any other purpose. Empty containers retain vapor and product residue. Observe all labeled safeguards until container is cleaned, reconditioned or destroyed. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller.

To clean the container before final disposal, empty the remaining contents from the container into formulation equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with pump for 2 minutes. Pour or pump rinsate into formulation equipment or rinsate collection system. Repeat this rinsing process two more times.

## CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

To the extent consistent with applicable law, upon purchase or use of this product, purchaser and user agree to the following terms:

**Warranty:** Alligare, LLC (the Company) warrants that this product conforms to the chemical description on the label in all material respects and is reasonably fit for the purpose referred to in the directions for use, subject to the exceptions noted below, which are beyond the Company's control. To the extent consistent with applicable law, the Company makes no other representation or warranty, express or implied, concerning the product, including no implied warranty of merchantability or fitness for a particular purpose. To the extent consistent with applicable law, no such warranty shall be implied by law, and no agent or representative is authorized to make any such warranty on the Company's behalf.

**Terms of Sale:** The Company's directions for use of this product must be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, and the manner of use or application (including failure to adhere to label directions), all of which are beyond the Company's control. To the extent consistent with applicable law, all such risks are assumed by the user.

**Limitation of Liability:** To the extent consistent with applicable law, the exclusive remedy against the Company for any cause of action relating to the handling or use of this product is a claim for damages, and in no event shall damages or any other recovery of any kind exceed the price of the product which caused the alleged loss, damage, injury or other claim. To the extent consistent with applicable law, under no circumstances shall the Company be liable for any special, indirect, incidental or consequential damages of any kind, including loss of profits or income. Some states do not allow the exclusion or limitation of incidental or consequential damages.

The Company and the seller offer this product, and the purchaser and user accept this product, subject to the foregoing warranty, terms of sale and limitation of liability, which may be varied or modified only by an agreement in writing signed on behalf of the Company by an authorized representative.

Amber, Beacon, and Tough are registered trademarks of Syngenta.

Accent, Express, Finesse, Glean, and Harmony are registered trademarks of E.I. duPont de Nemours & Co., Inc.

Bronate, Dakota, and Tiller are registered trademarks of Bayer CropScience.

Battalion, Bronco, Bullet, Harness, Landmaster, Lariat, Lasso, Partner, Permit, Ramrod, Roundup and Screen are registered trademarks of Monsanto Company.

Broadstrike, Curtail and Garlon are registered trademarks of Dow AgroSciences.

Clarity, Fallow Master, and Frontier are registered trademarks of BASF Corporation.

EPA 20160204